Keratoacanthoma Developing at the Site of Chemical Peel: A Rare Post-Procedure Complication

Erica M. Lin MD[1], Leticia Vidotti Storti MD[2], Raquel Steglich, MD[3], Carlos Gustavo Wambier, MD, PhD[4]

[1]University of Pittsburgh Department of Dermatology, Pittsburgh, PA, United States
[2]Brown University Department of Dermatology, Providence, RI, United States
[3]DermaClinic and Universidade da Região de Joinville, Joinville, SC, Brazil

Background

- Trauma is a recognized trigger for keratoacanthoma (KA) formation.
- To our knowledge, only one prior report has described KA development after a chemical peel using a non-phenol formulation; this is the first reported case following a phenol–croton oil peel.

Clinical Presentation

65 year old woman with no significant medical history

Pre-procedure

Baker-Gordon phenol peel had been well tolerated without intraoperative or immediate postoperative complications, excellent re-epithelialization at 5 weeks

2 months post-procedure

Discussion

- Injury-induced inflammation in chronically sun-damaged skin has been hypothesized to transiently activate MAPK signaling pathways, favoring the development of a KA over cutaneous squamous cell carcinoma after short-lived inflammatory stimuli, analogous to the paradoxical KA eruptions observed with BRAF inhibitor therapy.
- KA development has also been reported after aesthetic interventions, particularly laser resurfacing, in patients with actinically damaged skin. Although less common, other cosmetic treatments involving epidermal injury, such as chemical peels, may create a similar proinflammatory milieu conducive to KA formation.

Clinical Pearl

Dermatologists should maintain a high index of suspicion when a rapidly enlarging papule develops within a recently treated area. Early biopsy or excision enables definitive diagnosis and timely management, minimizing morbidity while preserving aesthetic outcomes.

References

Pattee SF, Silvis NG. Keratoacanthoma developing in sites of previous trauma: report of two cases and review of the literature. *J Am Acad Dermatol*. 2003;48(2 Suppl):S35-S38.

Cox S. Rapid development of keratoacanthomas after a body peel. *Dermatol Surg*. 2003;29(2):201-203.

Goldberg LH, Silapunt S, Beyrau KK, Peterson SR, Friedman PM, Alam M. *Keratoacanthoma as a postoperative complication of skin cancer excision*. *J Am Acad Dermatol*. 2004;50(5):753–758.

Su F, Viros A, Milagre C, et al. RAS mutations in cutaneous squamous cell carcinomas in patients treated with BRAF inhibitors. *N Engl J Med*. 2012;366(3):207-215. doi:10.1056/NEJMoa1105358

Witheiler DD, Chern PL, Quintana AT, Cockerell CJ. Multiple keratoacanthomas arising in sites of laser resurfacing. *Arch Dermatol*. 2002;138(4):483-485.